

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

THIS PAGE BLANK (USPTO)



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 35/14, 35/30, 38/18, C12N 5/08 // (A61K 35/14, 31:57) (A61K 38/18, 35:14)	A1	(11) International Publication Number: WO 98/41220 (43) International Publication Date: 24 September 1998 (24.09.98)
(21) International Application Number: PCT/IL98/00120 (22) International Filing Date: 13 March 1998 (13.03.98) (30) Priority Data: 08/818,818 14 March 1997 (14.03.97) US 09/041,280 11 March 1998 (11.03.98) US (71) Applicant (for all designated States except US): YEDA RE- SEARCH AND DEVELOPMENT CO. LTD. [IL/IL]; Weiz- mann Institute of Science, P.O. Box 95, 76100 Rehovot (IL). (72) Inventors; and (75) Inventors/Applicants (for US only): EISEN- BACH-SCHWARTZ, Michal [IL/IL]; Rupin Street 5, 76353 Rehovot (IL). SPIGLER, Orly [IL/IL]; Pinsker Street 11, 76308 Rehovot (IL). (74) Agent: BEN-AMI, Paulina; Yeda Research and Development Co. Ltd., Weizmann Institute of Science, P.O. Box 95, 76100 Rehovot (IL).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the</i> <i>claims and to be republished in the event of the receipt of</i> <i>amendments.</i>
(54) Title: MONONUCLEAR PHAGOCYTES AND THEIR USE TO PROMOTE AXONAL REGENERATION		
(57) Abstract Methods and compositions are disclosed for the use of allogeneic mononuclear phagocytes to promote axonal regeneration in the central nervous system of a mammal. In one embodiment, allogeneic mononuclear phagocytes are administered into the CNS at or near a site of injury or disease and the mammal is treated with at least one anti-inflammatory agent. In another embodiment, allogeneic mononuclear phagocytes are administered into the CNS at or near a site of injury or disease and one or more enumerated adjuvant factors (e.g. aFGF) are administered to the CNS; anti-inflammatory therapy of the mammal may optionally be added. In a further embodiment, allogeneic mononuclear phagocytes are first stimulated with one or more enumerated stimulatory factors and then administered into the CNS at or near a site of injury or disease. Methods for screening stimulatory tissue and cells are also disclosed.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**MONONUCLEAR PHAGOCYTES AND THEIR
USE TO PROMOTE AXONAL REGENERATION**

1. FIELD OF THE INVENTION

5 The present invention relates to compositions comprising mononuclear phagocytes, and to methods for using mononuclear phagocytes, to promote axonal regeneration in mammals affected by injury or disease of the central nervous system, as well as to compositions and methods for enhancing the
10 therapeutic capacity of mononuclear phagocytes to promote axonal regeneration. In particular, the invention relates to (a) pharmaceutical compositions comprising, and methods for administering, stimulated or non-stimulated allogeneic mononuclear phagocytes at or near a site of the mammalian
15 central nervous system affected by injury or disease to promote axonal regeneration, (b) compositions and methods for stimulating mononuclear phagocytes so as to enhance their capacity to promote axonal regeneration, and (c) methods for screening tissues, cells, proteins, peptides and other
20 biologically active agents for their ability to stimulate mononuclear phagocytes for promoting axonal regeneration.

2. BACKGROUND OF THE INVENTION

 Following axonal injury, neurons of the mammalian
25 central nervous system (CNS) have a poor capacity for axonal regeneration. By contrast, neurons of the mammalian peripheral nervous system (PNS) have a substantially greater capacity for axonal regeneration. See Schwartz et al., 1989, FASEB J. 3:2371-2378.

30 The difference between axonal regeneration in the CNS and PNS has been attributed to the cellular environment of the neurons rather than to the neurons themselves. Following neuronal injury, the Schwann cells that surround PNS neurons are modulated so as to become permissive or supportive for
35 axonal regeneration. By contrast, the astrocytes, oligodendrocytes and microglia that surround CNS neurons do not show such modulation and remain unsupportive or

inhibitory for axonal regeneration. See Schwartz et al., 1987, CRC Crit. Rev. Biochem. 22:89-110.

This lack of modulation has been correlated with differences in the post-injury inflammatory response. See 5 Perry and Brown, 1992, Bioessays 14:401-406; Lotan and Schwartz, 1994, FASEB J. 8:1026-1033. In particular, the accumulation of mononuclear phagocytes in response to CNS injury is delayed and limited in comparison with the response to injury in the PNS. This limited CNS mononuclear phagocyte 10 response may in turn lead to (1) inefficient removal of the myelin debris that reportedly inhibits axonal regeneration, and (2) suboptimal release of macrophage-derived cytokines that would promote modulation of astrocytes and oligodendrocytes so as to support axonal regeneration.

15 The above observations have prompted speculation that appropriate modulation of the macrophage response might promote axonal regeneration after CNS injury. In an *in vitro* system, David et al. showed that when cryostat sections of normal rat optic nerve are co-cultured with mononuclear 20 phagocytes derived from lesions of the rat CNS, the optic nerve sections show enhanced adhesiveness for embryonic chick dorsal root ganglion cells. David et al., 1990, Neuron 5:463-469. Conditioned medium from activated peritoneal macrophages was also effective in promoting adhesiveness of 25 optic nerve sections in this *in vitro* assay.

However, results derived from *in vivo* models of CNS injury have revealed that some interventions that enhance the macrophage response to CNS injury do not result in enhanced regeneration. For instance, local injection of either tumor 30 necrosis factor alpha (TNF- α) or colony stimulating factor-1 (CSF-1) enhanced the macrophage response to experimental optic nerve injury. However, only TNF- α , but not CSF-1, increased the permissiveness of the injured optic nerves for neuronal adhesion as assayed *in vitro*. Lotan et al., 1984, 35 Exp. Neurol. 126:284-290. It has been suggested as one possible explanation that "only appropriately stimulated

macrophages can influence neuronal regeneration." Schwartz et al., 1994, Progress Brain Res. 103:331-341, at 338.

In fact, contrary to the teaching of the present invention, other investigators have reported that mononuclear
5 phagocytes might exacerbate damage or limit recovery following CNS injury. Brain macrophages, when stimulated by cytokines, exhibit neurotoxic activity. Chamak et al., 1994, J. Neurosci. Res. 38:221-233. Pharmacological inhibition of mononuclear phagocyte function has been reported to promote
10 recovery in a rabbit model of spinal cord injury. Giulian and Robertson, 1990, Annals Neurol. 27:33-42. It has been suggested that macrophage-derived cytokines may promote formation of glial scars and thereby inhibit axonal regeneration. Khan and Wigley, 1994, NeuroReport 5:1381-
15 1385; Vick et al., 1992, J. Neurotrauma 9:S93-S103.

Citation or identification of any reference in Section 2 (or any other section) of this application shall not be construed as an admission that such reference is available as prior art to the present invention.

20

3. SUMMARY OF THE INVENTION

The present invention is directed to methods, and compositions, for use of allogeneic mononuclear phagocytes to promote axonal regeneration in the central nervous system of
25 a mammal. The allogeneic mononuclear phagocytes are administered into the CNS at or near a site of injury or disease.

In one embodiment, administration of allogeneic mononuclear phagocytes into the CNS at or near a site of
30 injury or disease is combined with administration to the mammal of at least one anti-inflammatory agent.

In another embodiment, administration of allogeneic mononuclear phagocytes into the CNS at or near a site of injury or disease is combined with administration to the CNS
35 of one or more adjuvant factors, for instance acidic fibroblast growth factor (aFGF), transforming growth factor-beta (TGF- β), interleukin 6 (IL-6), nerve growth factor

(NGF), neurotrophic factor 3 (NT-3), neurotrophic factor 4 (NT-4), neurotrophic factor 5 (NT-5) and brain-derived neurotrophic factor (BDNF). If desired, this can be combined with anti-inflammatory therapy of the mammal.

5 Allogeneic mononuclear phagocytes useful for the methods and compositions of the invention include, but are not limited to, allogeneic monocytes, macrophages and dendritic cells, and autologous monocytes, macrophages and dendritic cells.

10 The present invention further provides methods, and compositions, for stimulating allogeneic mononuclear phagocytes so as to enhance their capacity to promote axonal regeneration, and methods and compositions for use of stimulated allogeneic mononuclear phagocytes to promote
15 axonal regeneration in the central nervous system of a mammal. The mononuclear phagocytes are stimulated by culturing them in medium to which has been added one or more of the following: β -interferon (IFN- β), γ -interferon (IFN- γ), tumor necrosis factor α (TNF- α), interleukin 2 (IL-2),
20 interleukin 3 (IL-3), interleukin 4 (IL-4), interleukin 10 (IL-10), monocyte chemotactic and activating factor (MCAF), macrophage colony stimulating factor (M-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), colony stimulating factor 1 (CSF-1), lipid A, the tripeptide fMet-
25 Leu-Phe, muramyl dipeptide (MDP), the ionophore A23187, or vitamin D3 binding protein. A biologically active protein or peptide may be used in its native or recombinant form.

 The present invention further provides an assay for screening or identifying additional tissues, cells and
30 biologically active agents that are suitable for stimulating mononuclear phagocytes to enhance their capacity to promote axonal regeneration. According to this assay, mononuclear phagocytes are first cultured together with the tissue or cells to be tested, or in medium that has been conditioned by
35 the tissue or cells to be tested or in medium to which has been added the biologically active agent to be tested. The cultured mononuclear phagocytes are then assayed for nitric

oxide production, or both nitric oxide production and phagocytic activity. Mononuclear phagocytes with increased production of nitric oxide, or increased nitric oxide production and increased phagocytic activity, have an enhanced capacity to promote axonal regeneration.

4. BRIEF DESCRIPTION OF THE FIGURES

The present invention may be more fully understood by reference to the following detailed description of the invention, examples of specific embodiments of the invention and the appended figures in which:

FIGURE 1 illustrates axonal regeneration in transected optic nerves of rats as detected by retrograde transport of fluorescent dye to retinal ganglion cells (RGCs). See text, Section 6, for experimental details. Shortly after transection, 2 μ l of DCCM-1 medium were applied to the site of injury containing no cells (MED); 2.5×10^3 - 1×10^5 non-stimulated (NS) monocytes; 2.5×10^3 - 1×10^5 optic nerve-stimulated (OS) monocytes; or 2.5×10^3 - 1×10^5 sciatic nerve-stimulated (SS) monocytes. Open circles represent individual experimental animals. Solid circles represent animals that showed no labeled RGCs (numbering 7, 7 and 6 in the MED, NS and OS treatment groups respectively). Horizontal lines represent the median value of each treatment group.

FIGURE 2 illustrates axonal regeneration in transected optic nerves of rats as a function of the number and type of monocytes applied to the site of injury shortly after transection. See text, Section 6, for experimental details. At the time of transection, 2 μ l DCCM-1 medium were applied to the site of injury containing optic nerve-stimulated monocytes (OS) or sciatic nerve-stimulated monocytes (SS) at a total dose of 2.5×10^3 cells; 5×10^3 cells; 10^4 cells; or 10^5 cells.

FIGURE 3 (A-B) presents representative photomicrographs showing retrograde labeling of retinal ganglion cells in rats subjected to optic nerve transection followed by

administration of (A) 5×10^3 sciatic nerve-stimulated monocytes or (B) control medium. See text, Section 6, for experimental-details.

FIGURE 4 (A-E) presents representative photomicrographs showing anterograde labeling of optic nerve fibers in rats subjected to optic nerve transection followed by administration of sciatic nerve-stimulated monocytes (A-D) or control medium (E). See text, Section 6, for experimental details. FIGURE 4A is a low magnification view showing the point at which HRP was applied (H), the site of transection (ST) and the surrounding dura mater (DU). The bracketed region, distal to the site of transection, is shown at higher magnification in FIGURES 4B, 4C and 4D, in which growth cone-like structures (gc) are shown at the tips of the fibers.

FIGURE 5 illustrates axonal regeneration in transected optic nerves of rats after application to the site of injury of monocytes cultured with sciatic nerve for 2-17 hours. See text, Section 6, for experimental details. At the time of transection, $2 \mu\text{l}$ of DCCM-1 medium were applied to the site of injury containing 5×10^3 non-stimulated monocytes (NS) or 5×10^3 monocytes cultured with rat sciatic nerve for 2 hours (2h), 12 hours (12h) or 17 hours (17h).

FIGURE 6 illustrates axonal regeneration in transected optic nerves after administration, at the site of injury, of rat monocytes stimulated with mouse sciatic nerve or rat sciatic nerve. See text, Section 6, for experimental details. At the time of transection, $2 \mu\text{l}$ DCCM-1 medium were applied to the site of injury containing 5×10^3 monocytes cultured for 24 hours with either mouse sciatic nerve (MOUSE) or rat sciatic nerve (RAT).

FIGURE 7 illustrates the phagocytic activity of rat monocytes cultured for 2 hours with rat sciatic nerve. See text, Section 6, for experimental details. 2.5×10^5 rat monocytes were cultured in 1 ml DCCM-1 medium alone (CONTROL) or in 1 ml DCCM-1 medium with 2 segments of rat sciatic nerve (2SS) or with 4 segments of rat sciatic nerve (4SS). After 2

hours, the monocytes were exposed to fluorescent beads and cell-associated fluorescence was measured by flow cytometry.

FIGURE 8 illustrates the phagocytic activity of rat monocytes cultured for 24 hours with rat sciatic nerve. See 5 text, Section 6, for experimental details. 2.5×10^5 rat monocytes were cultured in 1 ml DCCM-1 medium alone (CONTROL) or in 1 ml DCCM-1 medium with 1 segment of rat sciatic nerve (1SS) or with 4 segments of rat sciatic nerve (4SS). After 16-24 hours, the monocytes were exposed to fluorescent beads 10 and cell-associated fluorescence was measured by flow cytometry.

FIGURE 9 illustrates the phagocytic activity of rat monocytes cultured for 2 hours with rat optic nerve. See 15 text, Section 6, for experimental details. 2.5×10^5 rat monocytes were cultured in 1 ml DCCM-1 medium alone (CONTROL) or in 1 ml DCCM-1 medium with 4 segments of rat optic nerve (4OS). After 2 hours, the monocytes were exposed to fluorescent beads and cell-associated fluorescence was measured by flow cytometry.

20 FIGURE 10 illustrates the phagocytic activity of rat monocytes cultured for 24 hours with rat optic nerve. See text, Section 6, for experimental details. 2.5×10^5 rat monocytes were cultured in 1 ml DCCM-1 medium alone (CONTROL) or in 1 ml DCCM-1 medium with 4 segments of rat optic nerve 25 (4OS). After 24 hours, the monocytes were exposed to fluorescent beads and cell-associated fluorescence was measured by flow cytometry.

FIGURE 11 illustrates the phagocytic activity of rat monocytes cultured overnight with rat sciatic nerve in the 30 presence of medium conditioned by rat optic nerve. 5×10^5 rat monocytes were cultured in 1 ml DCCM-1 medium with 6 segments of rat sciatic nerve with no further additions (0) or with the addition of optic nerve-conditioned medium at a total protein concentration of $0.1 \mu\text{g/ml}$ (0.1), $1.0 \mu\text{g/ml}$ 35 (1), or $10 \mu\text{g/ml}$ (10). After 24 hours, the monocytes were exposed to fluorescent beads and cell-associated fluorescence was measured by flow cytometry.

FIGURE 12 illustrates nitric oxide production by rat monocytes cultured for 24, 48, 72 or 96 hours with rat sciatic nerve or with rat optic nerve. See text, Section 6, for experimental details. 10^6 rat monocytes were cultured in 1 ml DCCM-1 medium alone (CONTROL), or in the same medium with 1 segment of rat sciatic nerve (1SS), with 1 segment of rat optic nerve (1OS), or with four segments of rat optic nerve (4OS). After 24, 48, 72 or 96 hours, the media were collected and the levels of nitric oxide were measured.

FIGURE 13 illustrates nitric oxide production by rat monocytes cultured for 72 hours with medium conditioned by rat sciatic nerve or rat optic nerve. See text, Section 6, for experimental details. 10^6 rat monocytes were cultured in 1 ml DCCM-1 medium with no further additions or with the addition of sciatic nerve-conditioned medium or optic nerve-conditioned medium at a total protein concentration of 10, 100, 200 or 300 $\mu\text{g/ml}$. After 72 hours, the media were collected and the levels of nitric oxide were measured.

FIGURE 14 illustrates axonal regeneration in transected optic nerves of rats following administration of optic nerve-stimulated monocytes combined with anti-inflammatory therapy. See text, Section 6, for experimental details. At the time of transection, 2 μl DCCM-1 medium were applied to the site of injury containing no cells or 5×10^3 sciatic nerve-stimulated rat monocytes. Concurrently, some of the rats received an intraperitoneal injection of 0.8 mg dexamethasone, producing the following treatment groups: no therapy (CONTROL), dexamethasone only (DEX), monocytes only (SS), and both dexamethasone and monocytes (DEX/SS).

FIGURE 15 illustrates recovery of voluntary motor function following administration of stimulated rat monocytes to rats that have undergone complete spinal cord transection. See text, Section 7, for experimental details. In FIGURE 15A, the solid line presents the BBB locomotor scores (mean \pm SEM) for 8 out of 12 animals that showed motor recovery after spinal cord transection and treatment with stimulated monocytes, and the broken line presents the BBB locomotor

scores of control animals following spinal cord transection. FIGURE 15B shows serial BBB locomotor scores for an individual animal subjected to spinal cord transection and treated with 4×10^5 stimulated monocytes plus aFGF.

5 FIGURE 16 presents low-power micrographs of transected spinal cord in rats treated with stimulated monocytes (A) or control medium (B), processed for immunohistochemical detection of GFAP (a) or neurofilament antigens (b). See text, Section 7, for experimental details. Each picture is a
10 montage of approximately 100 frames, each photographed at 10x magnification.

FIGURE 17 presents high-power micrographs of transected spinal cord in rats treated with stimulated monocytes (A) or control medium (B), processed for immunohistochemical
15 detection of neurofilament antigens (a) or GAP-43 (b). See text, Section 7, for experimental details. Bar: 2.5 μ m.

5. DETAILED DESCRIPTION OF THE INVENTION

5.1 MONONUCLEAR PHAGOCYTES

20 The present invention provides methods and compositions for use of allogeneic mononuclear phagocytes to promote axonal regeneration following injury or disease of the central nervous system (CNS). Allogeneic mononuclear phagocytes are introduced at or near the site of CNS injury
25 or disease.

As used herein, the term "mononuclear phagocytes" is intended to comprise, without limitation, monocytes obtained from central or peripheral blood, macrophages obtained from any site, including any tissue or cavity, macrophages derived
30 by culturing macrophage precursors obtained from bone marrow or blood, dendritic cells obtained from any site, including spleen, lymph node, skin and lymphatic fluid, and dendritic cells derived from culturing dendritic cell precursors obtained from bone marrow or blood.

35 Allogeneic mononuclear phagocytes can be obtained from the circulation or from any tissue in which they reside. Peripheral blood is an easily accessible ready source of

allogeneic monocytes and is used as a source according to a preferred embodiment of the invention. Especially preferred is the use of autologous monocytes purified from the peripheral blood of a subject to whom the therapeutic
5 preparation is intended to be administered.

Allogeneic mononuclear phagocytes from other sources are well known in the art and include, without limitation, macrophages obtained from serosal cavities such as the peritoneal or pleural cavity, alveolar macrophages, and
10 macrophages associated with other tissues (e.g. liver, spleen, thymus), where they may be known by various terms such as Kupffer cells (in the liver) and microglial cells (in the CNS). Allogeneic mononuclear phagocytes further include dendritic cells, which likewise may be known by various
15 terms, such as Langerhans cells (in the skin), veiled cells (in lymphatic fluid) and interdigitating cells (in lymph nodes). Additionally mononuclear phagocytes can be derived by culture from allogeneic brain-derived mixed glial cells or from allogeneic precursor cells, which may be obtained from
20 bone-marrow or blood. Preferably, the allogeneic mononuclear phagocytes are not microglia and are not derived by culture from brain-derived mixed glial cells.

In a preferred embodiment, cells other than mononuclear phagocytes are depleted from the cell population to be
25 administered. Enrichment techniques are well known to those skilled in the art and include, without limitation, elutriation; centrifugation through material of suitable density, such as a Percoll gradient (Colotta et al., 1983, J. Immunol. 132:936-944); selective adhesion on suitable
30 surfaces followed by removal at reduced temperature or at reduced concentrations of divalent cations (Rosen and Gordon, 1987, J. Exp. Med. 166:1685-1701), mechanical removal, or removal in the presence of lidocaine; and techniques for isolating dendritic cells from blood (O'Doherty et al., 1993,
35 J. Exp. Med. 178:1067-1078), bone marrow (Inaba et al., 1992, J. Exp. Med. 176:1693-1702) and lymphoid tissue (Macatonia et al., J. Exp. Med. 169:1255-1264). Preferably, at least 50%,

more preferably at least 70%, still more preferably at least 80%, and yet more preferably at least 90% of the cells are mononuclear phagocytes. Especially preferred is a substantially purified preparation of mononuclear phagocytes. 5 e.g. a preparation in which at least 95% of the cells are mononuclear phagocytes.

Once the mononuclear phagocytes are obtained they may be used therapeutically at any desired time, according to the needs of the patient. The mononuclear phagocytes may, if 10 desired, be cultured prior to administration in any suitable culture medium. Preferably, the mononuclear phagocytes are cultured in a vessel made from sterile material to which these cells show limited or no adherence. In a preferred embodiment, the mononuclear phagocytes are cultured in 15 sterile Teflon bags prior to administration.

As used herein, "stimulated" mononuclear phagocytes are mononuclear phagocytes with an enhanced capacity to promote axonal regeneration. Preferably, the capacity of the mononuclear phagocytes to promote axonal regeneration is 20 enhanced at least three-fold over non-stimulated mononuclear phagocytes, more preferably the capacity of the mononuclear phagocytes to promote axonal regeneration is enhanced at least 15-fold over non-stimulated mononuclear phagocytes. "Stimulatory" tissue, cells and biologically active agents 25 are tissue, cells and biologically active agents that, when cultured together with mononuclear phagocytes, enhance the capacity of the mononuclear phagocytes to promote axonal regeneration.

In a preferred embodiment, stimulatory tissue, cells or 30 at least one stimulatory biologically active agent is added to the culture in order to enhance the capacity of the mononuclear phagocytes to promote axonal regeneration. Preferably, one or more segments of a nerve, most preferably a peripheral nerve such as the sciatic nerve, are added to 35 the culture. A xenogeneic nerve is suitable for this purpose or, more preferably, an allogeneic or autologous nerve. If desired, a human nerve can be obtained from any available

human tissue, such as a human cadaver or a surgical specimen (e.g. an amputated limb). Alternatively other stimulatory tissue or cells are added to the culture. Dermis is suitable for this purpose and can be obtained, from a living donor or
5 a cadaver, by punch biopsy, by surgical resection, or by any other suitable technique. Especially preferred is skin obtained by punch biopsy, particularly skin obtained from a patient to whom the stimulated mononuclear phagocytes are intended to be administered. Synovial tissue, tendon sheath
10 and liver are also suitable for this purpose, as are other regenerating tissues. Additional stimulatory tissues and cells can be identified according to the assay described below. If desired, the stimulatory tissue or cells are homogenized before addition to the culture. As will be
15 evident to those skilled in the art, the stimulatory tissue or cell homogenate can be preserved, e.g. by cryopreservation, before use.

In an alternative embodiment, at least one stimulatory biologically active agent is added to the culture in order to
20 enhance the capacity of the mononuclear phagocytes to promote axonal regeneration. Neurotrophic factor 3 (NT3), nerve growth factor (NGF), brain-derived neurotrophic factor, β -interferon (IFN- β), γ -interferon (IFN- γ), tumor necrosis factor α (TNF- α), interleukin 2 (IL-2), interleukin 3 (IL-3),
25 interleukin 4 (IL-4), interleukin 10 (IL-10), monocyte chemotactic and activating factor (MCAF), macrophage colony stimulating factor (M-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), colony stimulating factor 1 (CSF-1), lipid A, fMet-Leu-Phe (fMLP), muramyl dipeptide
30 (MDP), the ionophore A23187, and vitamin D3 binding protein are suitable for this purpose either singly or in combination. Additional stimulatory biologically active agents (including additional stimulatory proteins and peptides) can be identified according to the assay described
35 below.

A biologically active protein or peptide may be used in its native or recombinant form, at a concentration (for each

protein or peptide) of 1 to 5000 ng/ml, more preferably 10 to 5000 ng/ml, still more preferably 100 to 2500 ng/ml, most preferably about 1000 ng/ml. In one embodiment, mononuclear phagocytes are stimulated by culturing them in medium to
5 which IL-4 or IL-10 (and more preferably both IL-4 and IL-10) have been added.

Preferably, the mononuclear phagocytes are cultured together with stimulatory tissue, stimulatory cells, homogenate of stimulatory tissue or stimulatory cells, or at
10 least one stimulatory biologically active agent for 24 hours. Shorter periods of culture, such as approximately 2 hours, are also effective, as are longer periods of culture, such as one or more weeks. In an alternative embodiment, stimulatory conditioned medium is prepared by incubating stimulatory
15 tissue or cells, preferably one or more segments of a nerve, most preferably a peripheral nerve such as the sciatic nerve, in any medium that is suitable for culturing mononuclear phagocytes. After removal of the tissue or cells, mononuclear phagocytes are cultured in the stimulatory
20 conditioned medium in order to enhance their capacity to promote axonal regeneration. After removal of the tissue or cells, the stimulatory conditioned medium can be stored and later used as desired for stimulating mononuclear phagocytes. Such stimulatory conditioned medium can be provided in the
25 form of a commercial kit. Preferably, the stimulatory conditioned medium is preserved during storage, for instance by refrigeration, whether as a liquid or as frozen medium. Alternatively, the stimulatory conditioned medium is lyophilized.

30 In a preferred embodiment, the mononuclear phagocytes are exposed to a tyrosine kinase inhibitor, such as tyrphostine, before, during, or after stimulation, so as to reduce or eliminate undesired mononuclear phagocyte activities, such as secretion of TNF- α .

35 As will be evident to those skilled in the art, the mononuclear phagocytes can be preserved, e.g. by cryopreservation, either before or after culture.

- Cryopreservation agents which can be used include but are not limited to dimethyl sulfoxide (DMSO) (Lovelock and Bishop, 1959, Nature 183:1394-1395; Ashwood-Smith, 1961, Nature 190:1204-1205), glycerol, polyvinylpyrrolidone
- 5 (Rinfret, 1960, Ann. N.Y. Acad. Sci. 85:576), polyethylene glycol (Sloviter and Ravdin, 1962, Nature 196:548), albumin, dextran, sucrose, ethylene glycol, i-erythritol, D-ribitol, D-mannitol (Rowe et al., 1962, Fed. Proc. 21:157), D-sorbitol, i-inositol, D-lactose, choline chloride (Bender
- 10 et al., 1960, J. Appl. Physiol. 15:520), amino acids (Phan The Tran and Bender, 1960, Exp. Cell Res. 20:651), methanol, acetamide, glycerol monoacetate (Lovelock, 1954, Biochem. J. 56:265), inorganic salts (Phan The Tran and Bender, 1960, Proc. Soc. Exp. Biol. Med. 104:388; Phan The Tran and Bender,
- 15 1961, in Radiobiology, Proceedings of the Third Australian Conference on Radiobiology, Ilbery, P.L.T., ed., Butterworth, London, p. 59), and DMSO combined with hydroxyethyl starch and human serum albumin (Zaroulis and Leiderman, 1980, Cryobiology 17:311-317).
- 20 A controlled cooling rate is critical. Different cryoprotective agents (Rapatz et al., 1968, Cryobiology 5(1):18-25) and different cell types have different optimal cooling rates. See, e.g., Rowe and Rinfret, 1962, Blood 20:636; Rowe, 1966, Cryobiology 3(1):12-18; Lewis et al.,
- 25 1967, Transfusion 7(1):17-32; and Mazur, 1970, Science 168:939-949 for effects of cooling velocity on survival of marrow-stem cells and on their transplantation potential. The heat of fusion phase where water turns to ice should be minimal. The cooling procedure can be carried out by use of,
- 30 e.g., a programmable freezing device or a methanol bath procedure.

Programmable freezing apparatuses allow determination of optimal cooling rates and facilitate standard reproducible cooling. Programmable controlled-rate freezers such as

35 Cryomed or Planar permit tuning of the freezing regimen to the desired cooling rate curve.

After thorough freezing, cells can be rapidly transferred to a long-term cryogenic storage vessel. In one embodiment, samples can be cryogenically stored in mechanical freezers, such as freezers that maintain a temperature of about -80°C or about -20°C . In a preferred embodiment, samples can be cryogenically stored in liquid nitrogen (-196°C) or its vapor. Such storage is greatly facilitated by the availability of highly efficient liquid nitrogen refrigerators, which resemble large Thermos containers with an extremely low vacuum and internal super insulation, such that heat leakage and nitrogen losses are kept to an absolute minimum.

Considerations and procedures for the manipulation, cryopreservation, and long term storage of hematopoietic stem cells, particularly from bone marrow or peripheral blood, are largely applicable to the mononuclear phagocytes of the invention. Such a discussion can be found, for example, in the following references, incorporated by reference herein: Gorin, 1986, Clinics in Haematology 15(1):19-48; Bone-Marrow Conservation, Culture and Transplantation, Proceedings of a Panel, Moscow, July 22-26, 1968, International Atomic Energy Agency, Vienna, pp. 107-186.

Other methods of cryopreservation of viable cells, or modifications thereof, are available and envisioned for use, e.g., cold metal-mirror techniques. See Livesey and Linner, 1987, Nature 327:255; Linner et al., 1986, J. Histochem. Cytochem. 34(9):1123-1135; see also U.S. Patent No. 4,199,022 by Senken et al., U.S. Patent No. 3,753,357 by Schwartz, U.S. Patent No. 4,559,298 by Fahy.

Frozen cells are preferably thawed quickly (e.g., in a water bath maintained at $37-41^{\circ}\text{C}$) and chilled immediately upon thawing. It may be desirable to treat the cells in order to prevent cellular clumping upon thawing. To prevent clumping, various procedures can be used, including but not limited to the addition before and/or after freezing of DNase (Spitzer et al., 1980, Cancer 45:3075-3085), low molecular weight dextran and citrate, hydroxyethyl starch (Stiff et

al., 1983, Cryobiology 20:17-24), or acid citrate dextrose (Zaroulis and Leiderman, 1980, Cryobiology 17:311-317), etc.

The cryoprotective agent, if toxic in humans, should be removed prior to therapeutic use of the thawed mononuclear
5 phagocytes. One way in which to remove the cryoprotective agent is by dilution to an insignificant concentration.

Once frozen mononuclear phagocytes have been thawed and recovered, they are used to promote axonal regeneration as described herein with respect to non-frozen mononuclear
10 phagocytes.

5.2 METHODS OF USE

In one embodiment of the present invention, the mononuclear phagocytes are suspended in a sterile
15 pharmaceutically acceptable carrier and administered into the CNS of a mammal, including a human subject, at or near a site of injury or disease. In a preferred embodiment, the pharmaceutically acceptable carrier is PBS, a culture medium, or any pharmaceutically acceptable fluid in which the
20 mononuclear phagocytes are suspended. However, alternative pharmaceutically acceptable carriers will readily be apparent to those skilled in the art.

If desired, treatment with mononuclear phagocytes may optionally be combined with local or systemic anti-
25 inflammatory therapy, for instance administration of (a) a steroid such dexamethasone or methylprednisolone, (b) a non-steroidal anti-inflammatory agent (NSAID) such as aspirin, indomethacin, ibuprofen, fenoprofen, ketoprofen or naproxen, or (c) an anti-inflammatory peptide, such as Thr-Lys-Pro
30 (TKP). The present invention encompasses the optional use of an anti-inflammatory agent at any dose that is effective in the subject to be treated. Such effective doses are well known to those skilled in the art and include, for example, standard-dose therapy, such as systemic methylprednisolone
35 100 mg daily for a human adult, and high-dose therapy, such as systemic methylprednisolone 1000 mg daily for a human adult.

In accordance with the present invention, treatment with mononuclear phagocytes may optionally be combined with concurrent administration to the CNS of one or more adjuvant factors. Adjuvant factors suitable for this purpose include 5 acidic fibroblast growth factor (aFGF); transforming growth factor-beta (TGF- β); interleukin 6 (IL-6); neurotrophic factors, e.g. nerve growth factor (NGF), neurotrophic factor 3 (NT-3), neurotrophic factor 4 (NT-4), neurotrophic factor 5 (NT-5), and brain-derived neurotrophic factor (BDNF); and the 10 neuronal cell adhesion molecule known as L1 (L1CAM) see Kallunki et al., 1997, J. Cell Biol. 138: 1343-1354. Acidic fibroblast growth factor (aFGF) is especially preferred. Each adjuvant factor can be administered at a dose of 6 to 10 ng/kg, either as a single dose or in repeated doses, e.g. at 15 weekly intervals. In one embodiment, one or more adjuvant factors are administered into the CNS at or near a site of disease or injury that is treated with mononuclear phagocytes, either together with, or shortly before or after administration of the mononuclear phagocytes. Alternatively, 20 one or more adjuvant factors are administered regionally, such as by intraventricular administration for treatment of the brain, by intrathecal administration for treatment of the spinal cord, or by intraocular administration for treatment of the retina or optic nerve. Both native and recombinant 25 adjuvant factors can be used. The present invention further encompasses combined treatment with (a) mononuclear phagocytes, (b) steroidal or non-steroidal anti-inflammatory therapy, and (c) one or more adjuvant factors.

In a preferred embodiment, the mononuclear phagocytes 30 are administered immediately following CNS injury and are introduced at the site of CNS injury, for example with a glass micropipette. However, the present invention encompasses administration of mononuclear phagocytes at any time (e.g. within a week, a fortnight, a month, 2 months, 3 35 months or 6 months) after the CNS sustains injury or disease, and encompasses introduction of the mononuclear phagocytes at

or near a site of CNS injury or disease by any neurosurgically suitable technique.

The compositions and methods of the present invention are useful for treating any injury or disease of the CNS that results in or is accompanied by axonal damage. The injury or disease may be situated in any portion of the CNS, including the brain, spinal cord, or optic nerve. One example of such injury or disease is trauma, including coup or countercoup injury, penetrating trauma, and trauma sustained during a neurosurgical operation or other procedure. Another example of such injury or disease is stroke, including hemorrhagic stroke and ischemic stroke. Yet another example of such injury or disease is optic nerve injury accompanying optic neuropathy or glaucoma. Still further examples of CNS injury or disease will be evident to those skilled in the art from this description and are encompassed by the present invention. The compositions and methods of the present invention are useful for treating CNS injury or disease that results in axonal damage whether or not the subject also suffers from other disease of the central or peripheral nervous system, such as neurological disease of genetic, metabolic, toxic, nutritional, infective or autoimmune origin.

The optimal dose of mononuclear phagocytes is proportional to the number of nerve fibers affected by CNS injury or disease at the site being treated. In a preferred embodiment, the dose ranges from about 2.5×10^3 to about 10^5 mononuclear phagocytes for treating a lesion affecting about 10^5 nerve fibers, such as a complete transection of a rat optic nerve, and ranges from about 2.5×10^4 to about 10^6 mononuclear phagocytes for treating a lesion affecting about 10^6 nerve fibers, such as a complete transection of a human optic nerve. More preferably, the dose ranges from about 10^4 to about 10^5 mononuclear phagocytes for treating a lesion affecting about 10^5 nerve fibers and ranges from about 10^5 to about 10^6 mononuclear phagocytes for treating a lesion

affecting about 10^6 nerve fibers. As will be evident to those of skill in the art, the dose of mononuclear phagocytes can be scaled up or down in proportion to the number of nerve fibers affected at the lesion or site of injury being
5 treated.

5.3 ASSAY FOR STIMULATORY TISSUES, CELLS AND BIOLOGICALLY ACTIVE AGENTS

The present invention provides an assay for identifying
10 stimulatory tissues and cells and stimulatory biologically active agents. Mononuclear phagocytes are cultured together with the tissue or cells to be tested, in medium conditioned by the tissue or cells to be tested, or in medium to which the biologically active agent or agents to be tested have
15 been added at various concentrations. Thereafter, the mononuclear phagocytes are assayed for phagocytic activity, or nitric oxide production. Mononuclear phagocytes with increased phagocytic activity or increased production of nitric oxide have an enhanced capacity to promote axonal
20 regeneration. In a preferred embodiment, both phagocytic activity and nitric oxide production are measured, and mononuclear phagocyte stimulation is detected by observing an increase in either of these activities, more preferably in both of these activities.

25 Preferably, the phagocytic capacity of the mononuclear phagocytes is increased by at least 10 percent, more preferably by at least 25 percent, still more preferably by at least 50 percent. Preferably, the nitric oxide production of the mononuclear phagocytes is increased by at least 50
30 percent, more preferably by at least 100 percent, and still more preferably by at least 200 percent.

In one embodiment, phagocytic activity is measured by contacting the mononuclear phagocytes with labeled particles and subsequently determining the amount of label associated
35 with the cells. A wide variety of particles can be used for this purpose, including without limitation latex or polystyrene beads and naturally occurring cells, such as red

blood cells, yeast and bacteria. Optionally, the particles can be opsonized, for instance with immunoglobulin or complement. The particles can be labeled with any suitable marker, including without limitation a fluorescent marker (such as fluorescein or rhodamine), a radioactive marker (such as a radioactive isotope of iodine, carbon or hydrogen), and an enzyme. Alternatively, the assay can be performed with unlabeled particles (e.g. red blood cells or yeast); the unlabeled particles are detected by any suitable method, such as microscopically, with or without staining. In a preferred embodiment, the mononuclear phagocytes are first contacted with fluorescent polystyrene beads; cell-associated fluorescence is subsequently measured by flow cytometry.

In one embodiment, nitric oxide production is measured by the Griess-reagent assay as described in Hibbs et al., 1987, Science 235:473-476, which is incorporated herein by reference. However, other assays for nitric oxide production may be used, as are known to those of skill in the art. See, e.g., Packer (ed.), 1996, Methods in Enzymology 268:58-247, which is incorporated herein by reference.

The assay of the present invention also provides a means of determining the period of culture required in order to stimulate the mononuclear phagocytes. Mononuclear phagocytes are cultured for various periods with stimulatory tissue or cells, in medium conditioned by stimulatory tissue or cells, or in medium to which at least one stimulatory biologically active agent has been added. Thereafter, the phagocytic activity or nitric oxide production of the mononuclear phagocytes, or both these properties, are measured. A period of culture sufficient to increase the phagocytic activity of the mononuclear phagocytes by at least 10 percent, preferably by at least 25 percent, more preferably by at least 50 percent, or sufficient to increase the nitric oxide production of the mononuclear phagocytes by at least 50 percent, preferably by at least 100 percent, more preferably

by at least 200 percent, is sufficient to stimulate their capacity to enhance axonal regeneration.

The following examples are presented for purposes of illustration only and are not intended to limit the scope of the invention in any way.

**6. EXAMPLE: USE OF MONOCYTES TO PROMOTE
AXONAL REGENERATION IN TRANSECTED OPTIC NERVE**

6.1 MATERIALS AND METHODS

6.1.1 ISOLATION AND CULTURE OF MONOCYTES

Peripheral blood was pooled from adult Sprague-Dawley (SPD) rats. Monocytes were isolated by fractionation on a one-step Percoll gradient as previously described. F. Colotta et al., 1984, J. Immunol. 132:936-944. The monocyte-enriched fraction was recovered from the Percoll interface, washed once with PBS to remove traces of Percoll, and resuspended at 1×10^6 cells/ml in DCCM-1 medium (Beit Ha'emek Ltd., Kibbutz Beit Ha'emek, Israel). The cells were cultured in Teflon bags at 37°C as previously described, Andreessen et al., 1983, J. Immunolog. Meth. 56:295-304, with 5% CO₂, as is conventional in the art. Usually, each bag received 10 ml containing 1×10^7 cells. For measurement of phagocytic activity or nitric oxide production, monocytes from SPD or Wistar rats were used, and were cultured in polypropylene tubes or in Teflon bags.

6.1.2 STIMULATION OF MONOCYTES

Non-stimulated monocytes (NS) were prepared by culturing isolated monocytes in a Teflon bag or polypropylene tubes, as described above, for 2-24 hours. Sciatic nerve-stimulated monocytes (SS) were prepared by culturing monocytes in a Teflon bag or polypropylene tubes for 2-24 hours together with at least one segment of a rat sciatic nerve. Optic nerve-stimulated monocytes (OS) were prepared by culturing monocytes in a Teflon bag or polypropylene tubes for 2-24 hours together with at least one segment of a rat optic nerve. Each nerve segment was 1.0 - 1.5 cm long in

experiments 6.2.1 and 6.2.2, and was 0.5 - 1.0 cm long in experiments 6.2.3 to 6.2.9; a constant ratio of 1 nerve segment to 5×10^6 cultured monocytes was used, except where otherwise noted.

5 After 2-24 hours in culture, monocytes were centrifuged for 3 minutes at 1000 x g, washed once with phosphate buffered saline (PBS), and resuspended in DCCM-1 medium at 1.25×10^6 - 5×10^6 cells/ml. The monocytes were 95% pure as determined by morphology and by immunocytochemistry with the
10 monoclonal antibody ED1 (Serotec, Oxford, England) as described. Hirschberg et al., 1994, J. Neuroimmunol. 50:9-16

Skin was also used to stimulate monocytes. In some experiments, 10^6 rat monocytes were cultured with a 1 cm x 1 cm square of skin obtained from germ-free rats by punch
15 biopsy. In other experiments, rat skin was cultured in protein-free medium to produce skin-conditioned medium containing skin-derived proteins; 10^6 rat monocytes were then cultured with skin-conditioned medium containing 200 μ g of protein.

20

6.1.3 OPTIC NERVE TRANSECTION

Anesthetized adult SPD rats, 8-9 weeks old, average mass 300 grams, were subjected to optic nerve transection as described. Eitan et al., 1994, Science 264:1764-1768. The
25 left optic nerve was exposed through a small opening in the meninges. A curved glass dissector with a 200 μ m tip and a smooth blunt edge was moved across the nerve to create a complete transection 2-3 mm distal to the optic globe, taking care not to damage the peripheral blood vessels. As used
30 herein, the term "distal" means away from the optic globe and towards the brain. Shortly after transection, 2 μ l of medium containing cultured monocytes or 2 μ l of medium alone were introduced at the site of injury by means of a curved glass micropipette with a 25 μ m lumen. The meningeal opening was
35 made about 200 μ m from the site of transection, in order to minimize leakage of cells from the site of application. In some experiments, dexamethasone (0.8 mg/rat) was administered

by intraperitoneal injection to some of the rats shortly after transection.

6.1.4 ASSAYS FOR AXONAL REGENERATION

5 6.1.4.1 RETROGRADE LABELING OF AXONS

Seven to eight weeks following transection, the lipophilic neurotracer dye, 4-(4-(didecylamino)styryl)-N-methylpyridinium iodide (4Di-10ASP) (Molecular Probes, Eugene, Oregon, USA) was applied to the injured optic nerve, 10 2 mm distal to the site of injury. One week after application of the dye, the retina was removed, prepared as a flattened whole mount in 4% paraformaldehyde solution, and examined by fluorescence microscopy to detect and count the number of labeled retinal ganglion cells (RGCs) in the entire 15 retina. Only axons that had regrown past the site of injury to the site at which dye was applied could take up the dye and transport it retrogradely to the retinal ganglion cells.

When applied to rat optic nerves that have not previously been transected, this procedure labels an average 20 of 21,623 RGCs per retina. The results for optic nerves that were subjected to transection are expressed as a percentage of this standard, to control for the efficiency of the 4Di-10ASP labeling technique.

25 6.1.4.2 ANTEROGRADE LABELING OF AXONS

Seven to eight weeks following transection, a fresh incision was made in the previously transected optic nerve 1 mm proximal to the site of transection. As used herein, "proximal" means towards the optic globe and away from the 30 brain. Horseradish peroxidase (HRP) (type VI-A, Sigma, Tel Aviv, Israel) was introduced through the incision by means of a sterile swab soaked in a 50% (w/v) solution of HRP in PBS. Eight to twelve hours after application of the HRP, the rats were perfused through the carotid artery with PBS followed by 35 4% paraformaldehyde in PBS as a fixative. The optic nerves were excised, 50 μ m longitudinal cryosections were taken and processed for visualization of HRP activity using

diaminobenzidine and cobalt intensification as described.
Lavie et al., 1992, Brain Res. 575:1-5.

6.1.5 ASSAY OF PHAGOCYTOTIC ACTIVITY

5 Monocytes from SPD or Wistar rats were suspended in
DCCM-1 medium (2.5×10^5 or 5×10^5 cells in 1 ml) and were
cultured without further additions or together with the
indicated number of syngeneic rat sciatic or optic nerve
segments or with the addition of medium conditioned by
10 syngeneic rat optic nerve at the indicated concentrations of
total protein. See Section 4 for details. To assay
phagocytic activity, a working solution of fluorescent
noncarboxylated microspheres ("FLUORESBRITE"TM, Polysciences,
Warrington, Pennsylvania, USA, Catalog. No. 17152) was
15 prepared by diluting 1 drop of a stock solution in 10 ml
DCCM-1 medium and adding this working solution to the
monocyte suspension at a further dilution of 1:100, after
removing the nerve segments. After three hours at 37°C, the
cells were washed once with DCCM-1 medium or with phosphate-
20 buffered saline, and cell-associated fluorescence was
measured by flow cytometry (FACS). In some experiments, 1 ml
ice-cold PBS was added prior to the washing step, to halt
phagocytosis.

6.1.6 ASSAY OF NITRIC OXIDE PRODUCTION

25 Monocytes from SPD or Wistar rats were suspended in
DCCM-1 medium (10^6 cells in 1 ml) and were cultured without
further additions or with the indicated number of syngeneic
rat sciatic or optic nerve segments or with the addition of
30 medium conditioned by syngeneic rat sciatic or optic nerve at
the indicated concentrations of total protein. See Section 4
for details. After the indicated time in culture, the nerve
segments (if any) were removed, the samples were centrifuged,
and the supernatants were collected. To assay nitric oxide
35 production, 100 μ l aliquots of supernatant were incubated
with 100 μ l of Griess reagent (1% sulfanilamide, 10% N-(1-
naphthyl)-ethylene diamine hydrochloride in 2.5% H_3PO_4) at

room temperature for 10 minutes. Colorimetry was performed at 550 nm with an ELISA reader and the amount of nitric oxide calculated according to a reference curve using sodium nitrite (Sigma, Israel) as standard. The reaction medium
5 (DCCM-1) was used as a blank. In control experiments, optic and sciatic nerve segments were found to produce negligible amounts of nitric oxide.

6.2 RESULTS

10 6.2.1 PROMOTION OF AXONAL REGENERATION BY STIMULATED AND NON-STIMULATED MONOCYTES

Rats were subjected to optic nerve transection and treated at the time of injury with control medium or with 2.5×10^3 - 1×10^5 non-stimulated (NS) monocytes, 2.5×10^3 - 1×10^5 sciatic nerve-stimulated (SS) monocytes, or 2.5×10^3 - 1×10^5 optic nerve-stimulated (OS) monocytes.
15

The number of labeled retinal ganglion cells (RGCs) in rats from each treatment group is shown in Figure 1 as a percentage of RGCs labeled in normal optic nerves. Rats
20 receiving no cells showed almost no labeling of RGCs. Rats receiving NS monocytes showed labeling of modest numbers of RGCs, while treatment with OS monocytes resulted in labeling of greater numbers of RGCs. In rats receiving SS monocytes, the median number of labeled RGCs was over 5-fold higher than
25 in the rats treated with OS monocytes, and was about 15-fold higher than in the rats treated with NS monocytes.

6.2.2 AXONAL REGENERATION AFTER TREATMENT WITH VARIOUS DOSES OF SCIATIC NERVE- OR OPTIC NERVE-STIMULATED MONOCYTES

30 To study regeneration as a function of the dose of monocytes administered, rats were subjected to optic nerve transection and treated at the time of injury with OS monocytes or SS monocytes at a total dose of 2.5×10^3 ; 5×10^3 ; 1×10^4 ; or 1×10^5 cells.

35 The average number of labeled retinal RGCs in each treatment group is shown in Figure 2 as a percentage of RGCs labeled in normal optic nerves. RGC labeling was highest

after treatment with 5×10^3 SS monocytes. Higher or lower doses of SS monocytes promoted axonal regeneration but were less effective. Treatment with OS monocytes similarly promoted axonal regeneration, though less effectively. The peak effect, with both OS and SS monocytes, occurred at a dose of 5×10^3 monocytes; at higher or lower doses the beneficial effect on axonal regeneration was less marked.

Representative fluorescence micrographs of labeled RGCs in retinas after treatment with SS monocytes or control medium are shown in Figure 3. The absence of labeled RGCs following treatment with control medium indicates that transection was complete and that the labeled RGCs represent regenerating axons that traversed the site of transection and not merely fibers that escaped the experimental injury.

The photomicrographs in Figure 4 further verify that regrowth has occurred. In nerves treated with control medium (E) no labeled fibers could be seen distal to the site of HRP application. In nerves treated with SS monocytes (A-D) labeled fibers were seen emerging from the proximal part of the nerve, crossing the site of transection (ST) and extending distally. Structures resembling growth cones (gc) were observed at the tips of these labeled fibers.

6.2.3 AXONAL REGENERATION AFTER TREATMENT WITH MONOCYTES STIMULATED WITH RAT SCIATIC NERVE SEGMENTS FOR VARIOUS INTERVALS

To study the capacity of monocytes to promote axonal regeneration after stimulation for various intervals with sciatic nerve segments, rats were subjected to optic nerve injury and treated at the time of injury with 5×10^3 monocytes cultured with rat sciatic nerve segments for two hours (2h), twelve hours (12h) or seventeen hours (17h). The number of labeled RGCs in individual rats from each treatment group is shown in Figure 5 as a percentage of RGCs labeled in normal optic nerves. Monocytes showed an enhanced capacity to promote axonal regeneration after culture with sciatic nerve segments for each interval tested.

6.2.4 AXONAL REGENERATION AFTER TREATMENT WITH MONOCYTES STIMULATED WITH RAT OR MOUSE SCIATIC NERVE SEGMENTS

To compare the ability of sciatic nerve segments derived from rat and mouse to stimulate the capacity of monocytes to promote axonal regeneration, rats were subjected to optic nerve transection and treated at the time of injury with 5×10^3 rat monocytes cultured for 24 hours either with 1-8 segments of rat sciatic nerve (RAT) or with 2-16 segments of mouse sciatic nerve (MOUSE). The number of labeled RGCs in individual rats from each treatment group is shown in Figure 6 as a percentage of RGCs labeled in normal optic nerves. Both rat and mouse sciatic nerve stimulated the capacity of monocytes to promote axonal regeneration.

6.2.5 PHAGOCYTTIC ACTIVITY OF MONOCYTES FOLLOWING CULTURE WITH SEGMENTS OF RAT SCIATIC NERVE

Rat monocytes were suspended at 2.5×10^5 cells in 1 ml DCCM-1 medium and were cultured for 2-24 hours without further additions (CONTROL), with 1 segment of rat sciatic nerve (1SS), with 2 segments of rat sciatic nerve (2SS), or with 4 segments of rat sciatic nerve (4SS).

The phagocytic activity of the 2SS and 4SS preparations after 2 hours in culture is shown in Figure 7 relative to the phagocytic activity of CONTROL monocytes. After culture for 2 hours with two segments of sciatic nerve, the monocytes showed increased phagocytic activity; after culture for 2 hours with four segments of sciatic nerve, the monocytes showed a greater increase in phagocytic activity.

The phagocytic activity of the 1SS and 4SS preparations after 24 hours in culture is shown in Figure 8 relative to the phagocytic activity of CONTROL monocytes. After culture for 24 hours with one segment of sciatic nerve, the monocytes showed increased phagocytic activity; after culture for 24 hours with four segments of sciatic nerve, the increase in phagocytic activity was even greater. The 4SS preparation

showed a greater increase in phagocytic activity after 24 hours than after 2 hours.

Addition of sciatic nerve-conditioned medium to the monocyte culture likewise increased the phagocytic activity of the monocytes (data not shown).

6.2.6 PHAGOCYTIC ACTIVITY OF MONOCYTES FOLLOWING CULTURE WITH SEGMENTS OF RAT OPTIC NERVE

Rat monocytes were suspended at 2.5×10^5 cells in 1 ml DCCM-1 medium and were cultured for 2-24 hours without further additions (CONTROL) or with 4 segments of rat optic nerve (40S). The phagocytic activity of the 40S preparations after 2 hours in culture is shown in Figure 9 relative to the phagocytic activity of CONTROL monocytes. After culture for 2 hours with four segments of optic nerve, the monocytes showed a decrease in phagocytic activity.

The phagocytic activity of the 40S preparations after 24 hours in culture is shown in Figure 10 relative to the phagocytic activity of CONTROL monocytes. After culture for 24 hours with four segments of optic nerve, the monocytes showed a decrease in phagocytic activity similar to that seen after 2 hours.

6.2.7 PHAGOCYTIC ACTIVITY OF MONOCYTES FOLLOWING CULTURE WITH SCIATIC NERVE SEGMENTS IN THE PRESENCE OF OPTIC NERVE-CONDITIONED MEDIUM

Optic nerve conditioned medium was prepared by culturing 10 segments of rat optic nerve for 2 hours in 1 ml DCCM-1 medium. While fresh DCCM-1 medium is protein-free, the optic nerve conditioned medium contained protein. Rat monocytes were suspended at 2.5×10^5 cells in 1 ml DCCM-1 medium and were cultured for 24 hours with 1-6 segments of rat sciatic nerve without further additions (0) or with optic nerve conditioned medium at a total protein concentration of 10 $\mu\text{g/ml}$ (10), 1 $\mu\text{g/ml}$ (1) or 0.1 $\mu\text{g/ml}$ (0.1).

Figure 11 presents the phagocytic activity of monocytes cultured with sciatic nerve in the presence of optic nerve

conditioned medium relative to the phagocytic activity of monocytes cultured with sciatic nerve in the absence of optic nerve conditioned medium. Addition of optic nerve conditioned medium attenuated the enhancement in phagocytic activity caused by culture with sciatic nerve. This attenuation was most marked in the preparation that received 0.1 µg/ml optic nerve conditioned medium. Similar results (not shown) were obtained when optic nerve segments were cultured in DCCM-1 medium for 8 hours and the resulting supernatants were dialyzed overnight at 4°C against PBS and subsequently stored at -20°C or -70°C.

6.2.8 NITRIC OXIDE PRODUCTION OF MONOCYTES CULTURED WITH SCIATIC NERVE, OPTIC NERVE, OR CONDITIONED MEDIUM

Rat monocytes were suspended at 10^6 cells in 1 ml DCCM-1 medium and were cultured for 24-96 hours without further additions (CONTROL), with 1 segment of rat sciatic nerve (1SS), or with 4 segments of rat optic nerve (4OS). The nitric oxide production of these preparations is shown in Figure 12. Monocytes cultured with sciatic nerve showed significantly increased production of nitric oxide, whereas optic nerve had no significant effect.

Figure 13 illustrates nitric oxide production of monocytes cultured for 72 hours with medium conditioned by rat sciatic nerve or rat optic nerve. Sciatic nerve-conditioned medium produced a statistically significant increase in nitric oxide production, whereas optic nerve-conditioned medium had no statistically significant effect. This result demonstrates that stimulation of mononuclear phagocytes by sciatic nerve is mediated by one or more soluble factors.

6.2.9 AXONAL REGENERATION AFTER TREATMENT WITH SCIATIC NERVE-STIMULATED MONOCYTES COMBINED WITH ANTI-INFLAMMATORY THERAPY

To study whether anti-inflammatory therapy prevents monocyte-mediated axonal regeneration, rats were subjected to

optic nerve transection. Control medium or sciatic nerve-stimulated monocytes were administered at the transection site shortly after injury, either without additional therapy or together with intraperitoneal administration of
5 dexamethasone. Eight weeks later, axonal regeneration was measured by retrograde labeling. As shown in Figure 14, rats receiving no therapy (CONTROL) or dexamethasone only (DEX) shown negligible regrowth, whereas sciatic nerve-stimulated monocytes promoted axonal regeneration, whether given alone
10 (SS) or concurrently with intraperitoneal dexamethasone (DEX/SS).

6.3 DISCUSSION

These examples demonstrate that monocytes administered
15 at a site of CNS injury promoted axonal regeneration. All monocytes tested were effective at promoting axonal regeneration. However, monocytes were stimulated (i.e., showed an enhanced capacity to promote axonal regeneration) by culture with a nerve segment, especially with a segment of
20 a peripheral nerve, e.g. sciatic nerve from rat or mouse. This stimulation was evident after all periods of culture tested, i.e. from 2-24 hours. For treating a total transection of a rat optic nerve, which contains about 10^5 nerve fibers, optimal results were obtained by administering
25 about 5×10^3 monocytes. However, every dose tested showed a beneficial effect on axonal regeneration.

These examples also demonstrate that monocytes show increased phagocytic activity and increased nitric oxide production after culture with one or more segments of sciatic
30 nerve or in sciatic nerve-conditioned medium. Thus, measurement of phagocytic activity, nitric oxide production or both these properties provides a rapid and efficient method of screening tissues and cells for their capacity to stimulate monocytes to promote axonal regeneration.

35

7. **EXAMPLE: USE OF MONOCYTES TO PROMOTE
AXONAL REGENERATION IN TRANSECTED SPINAL CORD**

7.1 **MATERIALS AND METHODS**

7.1.1 **ISOLATION AND STIMULATION OF MONOCYTES**

5 Peripheral blood from adult Sprague-Dawley (SPD) rats
was drawn into 10 ml syringes coated with heparin (5000 u/ml,
Calbiochem, La Jolla, CA), diluted with an equal volume of
PBS, and subjected to fractionation on a one-step gradient of
Percoll (1.077 g/ml, Pharmacia, Sweden) by centrifugation at
10 291 x g for 25 min at 30°C. See Colotta et al., 1984, J.
Immunol. 132: 936-944. The monocyte-enriched fraction was
recovered from the Percoll interface, washed once with PBS to
remove traces of Percoll, and resuspended at 1×10^6 cells/ml
in DCCM-1 medium (Beit Ha'emek Ltd., Kibbutz Beit Ha'emek,
15 Israel). The cells were incubated in polypropylene tubes or
Teflon bags at 37°C, 5% CO₂, with freshly excised segments of
rat sciatic nerve (0.5 to 1.0 cm long) for 2 to 24 hours (0.4
- 5.0×10^6 cells/nerve segment).

20 7.1.2 **SPINAL CORD TRANSECTION**

Male Sprague-Dawley rats (Hebrew University, Jerusalem,
Israel), 200-300 g, were anesthetized with ketamine 40 mg/kg
and xyline 100 mg/kg and incised dorsally to expose the T8-T9
vertebrae. The muscular insertions on the posterior and
25 transverse vertebral processes were dissected and cut with a
monopolar electrocautery device. T8 laminectomy was
performed with a bone rongeur, without contusive injury to
the underlying spinal cord. The spinal cord was transected
with microscissors, and any remaining fibers were cut with a
30 microknife. The underlying vertebral body was exposed
through a gap of approximately 3 mm between the cut ends of
the spinal cord. The exposed surface of the vertebral body
and the lateral recesses were checked under high
magnification to ensure that no fibers remained uncut.
35 During the procedure, bleeding was controlled with bipolar
electrocautery and by application of sterile gelfoam sponge
material (SPONGOSTAN™, Upjohn Co., Kalamazoo, MI).

Experimental protocols and procedures were in accordance with NIH guidelines for animal research.

Postoperatively, manual expression of the bladder was performed at least twice daily (and up to three times daily 5 in the first 48 hours), until autonomic bladder clearance developed at about the end of the second week. Some animals lost bladder automatism after the electrophysiological studies described below and required manual expression of the bladder until neurogenic control was recovered. The rats 10 were carefully monitored for evidence of urinary tract infection and any other signs of systemic disease. Trimethoprim and sulfamethoxazole (RESPRIM™, Teva Laboratories, Israel, 1 ml/day) was orally administered to each animal in the first postoperative week, and thereafter 15 to any animal showing hematuria. Daily inspection included examination of the laminectomy site for evidence of infection or serous collections, and assessment of the hind limbs for evidence of autophagia or pressure sores. Any severely ill rats were euthanized with an overdose of anesthesia. Such 20 rats, which never exceeded 10% of the number in any treatment group, were excluded for data analysis.

7.1.3 IMPLANTATION OF MONOCYTES

Syngeneic peripheral blood monocytes were purified by 25 one-step Percoll fractionation and co-cultured with segments of rat sciatic nerve, as described above. Prior to implantation, the sciatic nerve segments were removed and the cells were washed once and resuspended in fresh DCCM-1 medium and their viability determined.

30 In one group of animals, fibrin glue from a commercial kit (Octacol-FI5, OMRIX Biopharmaceuticals SA, Brussels, Belgium) was applied to the gap created between the cut ends of the spinal cord. First, 2.5 µl of the BAC component (which contains human fibrinogen 50 mg/ml and other human 35 plasma proteins, as well as tranexamic acid 92 mg/ml) was injected into the gap; then, 2.5 µl of the thrombin component (which contains human thrombin 1000 IU/ml and CaCl₂ 40 mM) was

injected into the gap. Following administration of fibrin glue, 5 μ l of monocyte suspension containing the indicated number of cultured monocytes (or control medium) were administered into the spinal cord parenchyma, distal (i.e. 5 caudal) to the site of injury, with a Hamilton syringe. In a second group of animals, fibrin glue was not used, and the monocyte suspension (or control medium) was injected partly into the gap and partly into the distal parenchyma. No significant difference was observed between these fibrin and 10 non-fibrin treatment groups, which were amalgamated for purposes of data analysis.

In some animals, aFGF (7.5 μ g/ml, 5 μ l/rat, Calbiochem Megapharm, Cat # 341580) was injected into the distal parenchyma. In all cases, the site of injury was covered 15 with a film of SPONGOSTAN™ and the wound was closed in layers.

7.1.4 EVALUATION OF MOTOR FUNCTION

Motor function was monitored using the open field 20 walking evaluation. Basso et al., 1995, J. Neurotrauma 12: 1-21; Basso et al., 1996, Exp. Neurol. 139: 244-256. Briefly, rats were placed in the middle of a circular enclosure of molded plastic (90 cm diameter, 7 cm wall height) with a smooth, non-slip floor. The rats were stimulated to provoke 25 continuous locomotion during the session, which lasted 4 minutes. A BBB locomotor score was assigned by observing movements involving the trunk, tail and hindlimbs and assigning a score ranging from 0 (no movement) to 22 (normal movement). All hindlimb movements were recorded except for 30 those that were obviously part of a reflex or that were elicited by a touch from the examiner. Special attention was given to hindlimb movements in the lower part of the open field score range.

To evaluate responses dependent on long ascending and 35 descending pathways, contact-placing reflexes were examined. Gale et al., 1985, Exp. Neurol. 88: 123-134; Kerasidis et al., 1987, J. Neurosci. Methods 20: 167-179. Briefly, this

response was elicited by light touching (hair bend) without proprioceptive stimulation, on the dorsal and lateral aspects of each foot. A positive contact-placing response involved flexion of the limb to clear the edge of the surface, and
5 subsequent extension and placement of the foot onto the surface for support.

7.1.5 ELECTROPHYSIOLOGICAL STUDIES

One week prior to spinal cord surgery, each rat
10 underwent implantation of screw electrodes extradurally over the sensorimotor cortex of each cerebral hemisphere. Simpson & Baskin, 1987, Neurosurgery 20: 131-137. Briefly, a hole was drilled on each side 1 mm lateral to the midline and 2 mm caudal to the coronal suture until the dura was exposed. A
15 3/16-inch screw was inserted (2.5 to 3 turns) into the hole. A thin layer of cyanoacrylate glue was applied to the surface of the skull and allowed to dry; the opening was then sealed with dental cement.

During electrophysiological recording, rats were
20 maintained under anesthesia (loading dose of ketamine 40 mg/kg and xylazine 10 mg/kg, administered ip and supplemented with one-third of the loading dose every 30 min). A ground needle electrode was inserted transdermally near the dorsum of the neck. Contralateral muscle motor evoked potentials
25 were elicited by stimulating the corresponding sensorimotor cortex with twin pulses of anodal stimulation from a Grass SD9 stimulator, applying 10 mA of constant current for 0.1 msec (with the cathodic electrode in the hard palate).

At least two motor-evoked potential traces averaging 50
30 sweeps were recorded from each muscle. A longitudinal skin incision was made along the anterior surface of each hindlimb. The aponeurotic layers were dissected to expose the gastrocnemius, tibialis anterior, quadriceps, adductors and biceps femoris. Monopolar needle electrodes were
35 inserted in the exposed muscles to capture the evoked EMG signals, which were amplified and filtered (Microelectrode AC Amplifier, model 1800, A-M Systems, Everett, WA; 100 Hz to 5

kHz bandpass); then digitized (LABVIEW® for Macintosh, National Instruments, Austin, TX); and then stored. After the procedure, the skin was sutured and prophylactic antibiotics were administered for the next few days.

5

7.1.6 IMMUNOHISTOCHEMICAL STUDIES

Spinal cord sections were indirectly immunostained for detection of glial fibrillary acidic protein (GFAP), neurofilament protein, or growth-associated protein (GAP-43).

- 10 Rats were perfused transcardially with saline and 4% paraformaldehyde/PBS solutions. The spinal cords were removed, postfixed in paraformaldehyde, and immersed overnight in 20% sucrose in PBS. Cryosections (20 µm) of macrophage-treated and control spinal cords were prepared,
- 15 placed on gelatin-coated slides, and dried at room temperature. Sections were fixed in absolute ethanol for 5 min at room temperature, washed several times in double-distilled water, and incubated for 5 min with 0.5% Tween-20 (Sigma, Israel) in PBS to enhance the permeability of the
- 20 tissue. Sections were incubated at 37°C with 5% bovine serum albumin in PBS for 30 min, then for 1-2 hours at room temperature with anti-GFAP antibody (Sigma, Israel; 1:100 dilution), anti-GAP-43 antibody (Boehringer-Mannheim, Germany; 1:100 dilution), or antibody raised against a
- 25 mixture of 68 kDa and 200 kDa neurofilament proteins (Novocastra Laboratories, UK; 1:50 dilution). After washing three times with 0.05% Tween-20/PBS, the sections were incubated for 30 min at room temperature with fluorescein-conjugated secondary antibodies (Jackson ImmunoResearch,
- 30 Jackson, PA; 1:100 dilution). After extensive washing, the sections were mounted in an antifading agent (1,4 diaza-bicyclo (2,2,2) octane; Sigma; 22 mM in PBS) and examined by fluorescence microscopy.

35

7.2 RESULTS

7.2.1 PROMOTION OF AXONAL REGENERATION IN THE TRANSECTED SPINAL CORD BY STIMULATED MONOCYTES

Rats were subjected to complete spinal cord transection and treated at the time of injury with (1) stimulated monocytes and aFGF; (2) stimulated monocytes alone; (3) (4) control medium and aFGF; and (4) control medium alone. The number of animals in each treatment group and the results obtained are summarized in Table 1.

Table 1. Functional recovery after spinal cord transection

TREATMENT		RESULTS	
Stimulated monocytes	aFGF	Number Operated	Number Recovered*
1-4 x 10 ⁵	+	12	9
1-4 x 10 ⁵	-	10	3
Nil (CONTROL)	+	10	0
Nil (CONTROL)	-	21	0

* Recovery is defined as open field locomotor score > 5

Animals were followed for 19 weeks after spinal cord transection. During the first 8 postoperative weeks, which included a brief initial period of spinal shock lasting about 2 days, all animals showed complete paralysis of the hindlimbs and there was no recovery of locomotor activity (i.e. no BBB locomotor score > 5) in any of the treatment or control groups.

After week 8, animals in the control groups (groups 3 and 4) showed either no movement of their hindlimbs or slight movement of one or two joints; the BBB locomotor score was mostly 2 or less, occasionally 4, and never exceeded 5. By contrast, significant recovery was seen in 12 of the 22 animals in the monocyte-treated groups (groups 1 and 2)

(ANOVA, $p < 0.001$). Recovery was manifested by extensive movement of all the joints of the hindlimbs, by plantar placement of the paws, and by weightbearing. Six of the treated animals attained a BBB locomotor score of 8 (i.e., sweeping movements with no weightbearing or plantar placement of the paw without weightbearing) and two treated animals attained a BBB locomotor score of 9 (i.e., plantar placement of the paw with weightbearing dorsal stepping and no plantar stepping, and a low-threshold contact placing response, which is considered to be a cortically-integrated reflex). Nine out of 12 animals treated with both monocytes and aFGF (group 1) showed recovery of motor activity, while 3 out of 10 animals treated with monocytes alone (group 2) showed motor recovery. Administering both aFGF and monocytes appeared to increase the proportion of recovered animals rather than the extent of improvement. However, animals that received aFGF alone, without monocytes, showed no motor recovery (i.e. BBB locomotor score did not exceed 5). Figure 1 (A and B) illustrates progressive recovery of motor function after spinal cord transection in rats treated with monocytes. Additional experiments (data not shown) suggest that the added benefit of aFGF is less marked when animals are treated with greater numbers of monocytes.

In a further treatment group (not shown) fibrin glue was placed in the gap between the cut ends of spinal cord; monocytes and aFGF were then administered into the fibrin (and not into the distal parenchyma). No recovery was seen in this treatment group, suggesting that recovery depends on an adequate number of monocytes being accessible to the transected nerve at the site of the lesion and distal to the site of injury.

Electromyographic studies confirmed that reinnervation of hindlimb muscles occurred in monocyte-treated rats. In normal rats, unipolar stimulation of the hindlimb area of the sensorimotor cortex (a) evokes a late hindlimb electromyographic response (latency 20-30 msec, threshold response at 3-4 mA) and (b) often evokes an early hindlimb

electromyographic response (latency 8-10 msec, threshold response at 8 mA). The late EMG response has been related to the corticospinal tract in the rat, on the basis of its conductive properties and lower threshold. Kalderon & Fuks, 5 1996, Proc. Nat'l Acad. Sci. USA 93: 11185-11190.

In the present study, no late EMG response was seen in any rat (including control and experimental groups) within 2 weeks after spinal cord transection. In control animals examined 12-14 weeks after transection (n=30), no late EMG 10 response was detected in even the most proximal muscle, the quadriceps. By contrast, muscles of monocyte-treated rats (groups 1 and 2) showed varying degrees of recovery of the late EMG response. Complete correlation was observed between the behavioral scores and the electromyographic observations. 15 All animals that showed voluntary locomotor recovery also showed a positive EMG response. Moreover, the locomotor and EMG observations correlated in terms of extent of recovery and bilaterality.

Immunohistochemical studies confirmed the presence of 20 regrowing fibers across the lesioned site. Regrowing fibers were detected using antibodies against the growth-associated protein, GAP-43. Adjacent serial sections of the same tissues were stained with anti-neurofilament antibodies (to detect nerve fibers) and with anti-GFAP antibodies (to detect 25 astrocytes). As shown in Figures 2 and 3, the histological observations correlated with the electrophysiological and behavioral results. As seen in Fig. 2 (Aa, Ba), the site of injury was delineated by GFAP staining. See Blaugrund et al., 1993, J. Comp. Neurol. 330: 105-112). In control animals, 30 the site of the lesion was devoid of neurofilament antigens and GAP-43. See Fig. 2 (Bb) & Fig. 3 (Ba, Bb) In monocyte-treated animals with hindlimb EMG responses, by contrast, the site of injury (delineated by GFAP staining) showed intense staining for neurofilament antigens and GAP-43, see Fig. 2 35 (Ab) & Fig. 3 (Aa, Ab). These results demonstrate that physiological recovery was associated with regrowth of nerve fibers across the site of the lesion. In monocyte-treated

animals that did not show recovery, the lesion site did not stain for neurofilament antigens or GAP-43 (not shown).

7.3 DISCUSSION

5 This example demonstrates that treatment with monocytes led to axonal regeneration, including functional recovery, in rats after complete transection of the spinal cord. Recovery was assessed by voluntary hindlimb motor function, scored in a open field, and by cortically evoked hind limb muscle
10 activity, detected by electromyography. There was excellent correlation between these distinct modes of assessing functional recovery. The motor recovery observed in the present study (values of 8-9 on the 21-point open field locomotor test scale) corresponds to a value of 2 to 3 on the
15 5-point open field test scale. See Cheng et al., 1996, Science 273: 510-513; Young, 1996, Science 273: 451. The beneficial effect of administering monocytes was enhanced by concurrent treatment with aFGF, although the added benefit of aFGF was less marked upon administration of higher numbers of
20 monocytes. When the monocytes were embedded in fibrin glue, rather than administered in a fluid suspension, no benefit was seen.

 The present invention is not to be limited in scope by
25 the exemplified embodiments, which are intended as illustrations of single aspects of the invention. Indeed, various modifications of the invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and
30 accompanying drawings. Such modifications are intended to fall within the scope of the appended claims.

 All publications cited herein are incorporated by reference in their entirety.

WHAT IS CLAIMED IS:

1. A method of promoting axonal regeneration in the central nervous system (CNS) of a mammal, comprising:
 - 5 (a) administering allogeneic mononuclear phagocytes into the CNS at or near a site of injury or disease; and
 - (b) administering to the mammal at least one anti-inflammatory agent.
- 10 2. A method of promoting axonal regeneration in the central nervous system (CNS) of a mammal, comprising:
 - (a) administering allogeneic mononuclear phagocytes into the CNS at or near a site of injury or
 - 15 disease; and
 - (b) administering to the CNS at least one of the following adjuvant factors: acidic fibroblast growth factor (aFGF), transforming growth factor-beta (TGF- β), interleukin 6 (IL-6), nerve growth
 - 20 factor (NGF), neurotrophic factor 3 (NT-3), neurotrophic factor 4 (NT-4), neurotrophic factor 5 (NT-5) and brain-derived neurotrophic factor (BDNF).
- 25 3. The method according to claim 2, wherein step (b) comprises administering acidic fibroblast growth factor (aFGF) to the CNS.
4. The method according to claim 3, wherein said aFGF is
- 30 administered at or near said site of injury or disease.
5. The method according to any one of claims 2 to 4, further comprising administering to the mammal at least one anti-inflammatory agent.

35

6. The method according to claim 1 or 5, wherein said anti-inflammatory agent is a steroid, a non-steroidal anti-inflammatory agent (NSAID), or the tripeptide Thr-Lys-Pro.

5 7. The method according to any one of claims 1 to 6, wherein said allogeneic mononuclear phagocytes are stimulated allogeneic mononuclear phagocytes.

8. The method according to claim 7, in which said
10 stimulated allogeneic mononuclear phagocytes have been produced by culturing allogeneic mononuclear phagocytes together with skin or with medium conditioned by skin.

9. The method according to claim 8, in which said skin is
15 autologous skin.

10. The method according to claim 7, in which said said stimulated allogeneic mononuclear phagocytes have been produced by culturing allogeneic mononuclear phagocytes
20 together with at least one nerve segment or with medium conditioned by at least one nerve segment.

11. The method according to claim 10, in which said nerve segment is a segment of a peripheral nerve.

25

12. The method according to claim 11, in which said nerve segment is a segment of an allogeneic peripheral nerve.

13. The method according to claim 7, in which said
30 stimulated allogeneic mononuclear phagocytes have been produced by culturing allogeneic mononuclear phagocytes in medium to which has been added β -interferon (IFN- β), γ -interferon (IFN- γ), tumor necrosis factor α (TNF- α), interleukin 2 (IL-2), interleukin 3 (IL-3), interleukin 4
35 (IL-4), interleukin 10 (IL-10), monocyte chemotactic and activating factor (MCAF), macrophage colony stimulating factor (M-CSF), granulocyte-macrophage colony stimulating

factor (GM-CSF), colony stimulating factor 1 (CSF-1),
neurotrophic factor 3 (NT-3), nerve growth factor (NGF),
brain-derived neurotrophic factor (BDNF), lipid A, the
tripeptide fMet-Leu-Phe, muramyl dipeptide (MDP), the
5 ionophore A23187, or vitamin D3 binding protein.

14. The method according to claim 13, in which said
stimulated allogeneic mononuclear phagocytes have been
produced by culturing allogeneic mononuclear phagocytes in
10 medium to which has been added IL-4, IL-10, or both IL-4 and
IL-10.

15. A method of promoting axonal regeneration in the central
nervous system (CNS) of a mammal, comprising: administering
15 the CNS, at or near a site of injury or disease, allogeneic
mononuclear phagocytes that have been cultured in medium to
which has been added β -interferon (IFN- β), γ -interferon (IFN- γ), tumor necrosis factor α (TNF- α), interleukin 2 (IL-2),
interleukin 3 (IL-3), interleukin 4 (IL-4), interleukin 10
20 (IL-10), monocyte chemotactic and activating factor (MCAF),
macrophage colony stimulating factor (M-CSF), granulocyte-
macrophage colony stimulating factor (GM-CSF), colony
stimulating factor 1 (CSF-1), lipid A, the tripeptide fMet-
Leu-Phe, muramyl dipeptide (MDP), the ionophore A23187, or
25 vitamin D3 binding protein.

16. The method according to any one of claims 1 to 15, in
which said allogeneic mononuclear phagocytes are autologous
mononuclear phagocytes.

30

17. The method according to any one of claims 1 to 16, in
which said mononuclear phagocytes are monocytes.

18. The method according to any one of claims 1 to 16, in
35 which said mononuclear phagocytes are macrophages.

19. The method according to claim 18, in which said mononuclear phagocytes are macrophages obtained from a serosal cavity, alveolar macrophages, macrophages obtained from the liver, spleen or thymus, or macrophages derived from
5 culturing macrophage precursors obtained from bone marrow or from blood.

20. The method according to any one of claims 1 to 16, in which said mononuclear phagocytes are not microglia and are
10 not derived by culture from brain-derived mixed glial cells.

21. The method according to any one of claims 1 to 16, in which said mononuclear phagocytes are dendritic cells.

15 22. The method according to any one of claims 1 to 21, in which said mammal is a human being.

23. The method according to any one of claims 1 to 22, in which said allogeneic mononuclear phagocytes are administered
20 into the spinal cord.

24. A pharmaceutical composition for administration to a mammal, said pharmaceutical composition comprising:

- (a) a pharmaceutically acceptable carrier; and
- 25 (b) mammalian mononuclear phagocytes that have been cultured in medium to which has been added β -interferon (IFN- β), γ -interferon (IFN- γ), tumor necrosis factor α (TNF- α), interleukin 2 (IL-2), interleukin 3 (IL-3), interleukin 4 (IL-4),
30 monocyte chemotactic and activating factor (MCAF), macrophage colony stimulating factor (M-CSF), granulocyte-macrophage colony stimulating factor (GM-CSF), colony stimulating factor 1 (CSF-1), lipid A, the tripeptide fMet-Leu-Phe, muramyl
35 dipeptide (MDP), the ionophore A23187, or vitamin D3 binding protein.

25. The pharmaceutical composition according to claim 24, in which said mammalian mononuclear phagocytes are human mononuclear phagocytes.
- 5 26. The pharmaceutical composition according to claim 24 or 25, in which said mononuclear phagocytes are monocytes.
27. The pharmaceutical composition according to claim 24 or 25, in which said mononuclear phagocytes are macrophages.
- 10 28. The pharmaceutical composition according to claim 24 or 25, in which said mononuclear phagocytes are dendritic cells.
29. The pharmaceutical composition according to any one of 15 claims 24 to 28, in which said said mammalian mononuclear phagocytes are human monocytes, macrophages or dendritic cells.
30. The pharmaceutical composition according to any one of 20 claims 24 to 29, for promoting axonal regeneration in the central nervous system (CNS) of a mammal.
31. The pharmaceutical composition according to claim 30, for administration into the CNS at or near a site of injury 25 or disease.
32. A method of detecting stimulatory activity comprising:
- 30 (a) culturing mononuclear phagocytes together with at least one tissue or with at least one type of cell, or with medium to which has been added at least one biologically active agent; and
- (b) after the step of culturing, measuring the nitric oxide production or both the phagocytic activity and the nitric oxide production of said mononuclear 35 phagocytes.

33. A method of preparing stimulated mononuclear phagocytes, comprising:

- 5 (a) culturing a first preparation of mononuclear phagocytes together with at least one tissue or with at least one cell type, or with medium conditioned by at least one tissue or at least one cell type, or with medium to which has been added at least one biologically active agent;
- 10 (b) after the step of culturing, measuring the nitric oxide production or both the phagocytic activity and the nitric oxide production of said first preparation of mononuclear phagocytes to identify a stimulatory tissue, stimulatory cell type or stimulatory biologically active agent; and
- 15 (c) culturing a second preparation of mononuclear phagocytes together with said stimulatory tissue or stimulatory cell type, with medium conditioned by said stimulatory tissue or stimulatory cell type, or with medium to which has been added said
- 20 stimulatory biologically active agent.

34. A preparation of stimulated mononuclear phagocytes produced by the method of claim 33.

25 35. A method of promoting axonal regeneration in the central nervous system (CNS) of a mammal, comprising:

- 30 (a) culturing a first preparation of allogeneic mononuclear phagocytes together with at least one tissue or with at least one cell type, or with medium conditioned by at least one tissue or at least one cell type, or with medium to which has been added at least one biologically active agent;
- (b) after the step of culturing, measuring the nitric oxide production or both the phagocytic activity
- 35 and the nitric oxide production of said first preparation of allogeneic mononuclear phagocytes to

identify a stimulatory tissue, stimulatory cell type or stimulatory biologically active agent;

- (c) culturing a second preparation of allogeneic mononuclear phagocytes together with said stimulatory tissue or stimulatory cell type, with medium conditioned by said stimulatory tissue or stimulatory cell type, or with medium to which has been added said stimulatory biologically active agent, so as to produce a preparation of stimulated mononuclear phagocytes; and
- (d) administering at least a portion of said preparation of stimulated mononuclear phagocytes into the CNS at or near a site of injury or disease.

1/18




FIG.1

2/18




FIG.2

3/18

FIG.3A



50μm

FIG.3B

4/18



FIG. 4B




FIG. 4C



FIG. 4D



FIG. 4E



5/18




FIG.5

6/18




FIG.6

7/18




FIG.7

8/18




FIG.8

9/18




FIG.9

10/18




FIG.10

11/18




FIG.11

12/18




Fig.12

13/18




Fig.13

15/18




FIG. 15A

16/18




FIG. 15B

17/18

Fig.16 A




Fig.16 B




Fig.16

18/18




Fig.17A

Fig.17B

INTERNATIONAL SEARCH REPORT

Intern: al Application No

PCT/IL 98/00120

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 A61K35/14 A61K35/30 A61K38/18 C12N5/08 //(A61K35/14,
31:57), (A61K38/18, 35:14)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	LAZAROV-SPIEGLER O ET AL: "Transplantation of activated macrophages overcomes central nervous system regrowth failure." FASEB JOURNAL, (1996 SEP) 10 (11) 1296-302, XP002072904 United States see the whole document ---	1-35
A	LOTAN M ET AL: "Cytokines modulate the inflammatory response and change permissiveness to neuronal adhesion in injured mammalian central nervous system." EXPERIMENTAL NEUROLOGY, (1994 APR) 126 (2) 284-90, XP002072905 United States see the whole document ---	1-35
-/--		



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

28 July 1998

Date of mailing of the international search report

12/08/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Moreau, J

INTERNATIONAL SEARCH REPORT

Internal Application No

PCT/IL 98/00120

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	CHAMAK B ET AL: "Brain macrophages stimulate neurite growth and regeneration by secreting thrombospondin." JOURNAL OF NEUROSCIENCE RESEARCH, (1994 JUN 1) 38 (2) 221-33, XP002072906 United States cited in the application see the whole document ---	1-35
P,X	LAZAROV-SPIEGLER O ET AL: "Transplantation of activated macrophages induces axonal regrowth following injury." 27TH ANNUAL MEETING OF THE SOCIETY FOR NEUROSCIENCE, NEW ORLEANS, LOUISIANA, USA, OCTOBER 25-30, 1997. SOCIETY FOR NEUROSCIENCE ABSTRACTS 23 (1-2). 1997. 1995, XP002072907 see the whole document ---	1-35
P,X	WO 97 09885 A (YEDA RESEARCH AND DEVELOPMENT CO.) 20 March 1997 see the whole document ---	1-35
P,X	WO 97 26901 A (YEDA RESEARCH AND DEVELOPMENT CO.) 31 July 1997 see the whole document -----	1-35

INTERNATIONAL SEARCH REPORT

Int. ational application No.

PCT/IL 98/00120

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 1-23 and 35
are directed to a method of treatment of the human/animal
body, the search has been carried out and based on the alleged
effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims: it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IL 98/00120

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9709885 A	20-03-1997	AU 7157296 A	01-04-1997
WO 9726901 A	31-07-1997	AU 1554197 A	20-08-1997